

Pr
Elin


**BREAST
CANCER
FUND**

What We Do | Events | Action Center | Giving | News | Resources | About Us

LOG IN | MEDIA CENTER

SEARCH

E-NEWSLETTER SIGN UP


IN THIS SECTION

Advocacy
Education & Outreach

Events
BCF Blog

Home > What We Do > Education & Outreach > Breast Cancer Fact Sheets and Resources > Chemical Fact Sheet: Parabens

Parabens and Breast Cancer

Parabens are synthetic chemicals used as a preservative to inhibit the growth of bacteria, yeasts, and molds. Parabens are known to disrupt endocrine (hormone) function. More than 12 research studies (1) show parabens to have estrogenic activity in animals and in tissue culture. Recent research detected five types of intact parabens in human breast tumors (2). Although not a conclusive link between exposure to parabens and breast cancer, this new research signals the need for a precautionary approach to the manufacture and use of these compounds.

The international research community is beginning to question the safety of parabens in consumer products, based on new evidence of their endocrine disrupting effects. These ubiquitous chemicals are used as preservatives in a host of consumer products: food, pharmaceuticals and personal care products, including shampoos and conditioners, sunscreens, and deodorants. Since 2000, 13 research studies have shown that various types of parabens (methyl, ethyl, propyl, benzyl and benzyl) act like estrogen in animals and in tissue culture. However, the estrogenic activity occurs only when parabens are applied to the skin, not when ingested.

The estrogen/breast cancer connection is indisputable, which raises concerns about chemicals such as parabens that behave like estrogen. These concerns include:

- The widespread use of parabens in countless consumer products makes them ubiquitous in the environment and in our bodies through multiple routes of exposure. A report published in the Journal of the American College of Toxicology in 1984 (3) estimated that parabens could be found in over 13,000 cosmetics products.
- Parabens are just one type of xenoestrogen to which women are exposed (xenoestrogens are synthetic agents that mimic the actions of estrogen). Consequently, another cause of concern is both the cumulative effect and the interaction of parabens with other xenoestrogens, and with the body's own estrogens – all of which affect endocrine function.
- A 2004 UK study detected traces of five parabens in the breast cancer tumors of 19 out of 20 women studied (4). This small study does not prove a causal relationship between parabens and breast cancer, but it is important because it detected the presence of intact parabens -- unaltered by the body's metabolism -- an indication of the chemicals' ability to penetrate skin and remain in breast tissue.

Daily use of personal care products – especially deodorants and anti-perspirants -- can result in direct, chronic exposure to parabens. A 2003 study published in the European Journal of Cancer Prevention (5) reported, "Frequency and earlier onset of antiperspirant/deodorant usage with underarm shaving were associated with an earlier age of breast cancer diagnosis."

The U.K. study highlights the need for more research on the potential link between products containing parabens and increased breast cancer risk (6). The authors also point out that prenatal exposure to parabens, and the potential effect on adult cancer risk, needs to be studied. Animal studies indicate that parabens also may affect development of the male reproductive system (7).

While research continues, however, the accumulated evidence makes a compelling case for taking a precautionary approach, individually and collectively, to the manufacture and use of parabens.

Footnotes

1. Blair, RM, Fang H, Branham WS, Hass BS, Dial SL, Moland CL, Tong W, Shi L, Perkins R, Sheehan DM (2000). The estrogen receptor relative binding affinities of 188 natural and xenochemicals: Structural diversity of ligands. Toxicological Sciences 54:138-153.

Byford JR, Shaw LE, Drew MGB, Pope GS, Sauer MJ, Darbre PD (2002). Oestrogenic activity of parabens in MCF7 human breast cancer cells. Journal of

RECENT PRESS COVERAGE

March 19, 2007, Rose Sheet: Campaign For Safe Cosmetics Unimpressed By CTFA Consumer Commitments

March 18, 2007, Contra Costa Times: Mount Shasta climb to aid cancer research

February 20, 2007, Seattle Post-Intelligencer: State mulls cosmetic safety law

February 16, 2007, Pacific Sun: Girl, you'll be a woman soon

February 15, 2007, New York Times: Should you trust your makeup?

[See All Press Coverage](#)

Steroid Biochemistry and Molecular Biology 80:49-60.

Darbre PD (2003). Underarm cosmetics and breast cancer. *Journal of Applied Toxicology* 23:89-95.

Darbre PD, Byford JR, Shaw LE, Hall S, Coldham NG, Pope GS, Sauer MJ (2003). Estrogenic activity of benzylparaben. *Journal of Applied Toxicology* 23:43-51.

Darbre PD, Byford JR, Shaw LE, Horton RA, Pope GS, Sauer M (2002). Oestrogenic activity of isobutylparaben in vitro and in vivo. *Journal of Applied Toxicology* 22:219-226.

Inui M, Tetsuya A, Takenaka S, Inui H, Nakazawa M, Ueda M, Watanabe H, Mori C, Iguchi T, Miyatake K (2003). Effects of UV screens and preservatives on vitellogenin and choriogenin production in male medaka (*Oryzias latipes*). *Toxicology* 194:43-50.

Jin-Sung P, Lee BJ, Kang KS, Tai JH, Cho JJ, Cho MH, Inoue T, Lee YS (2000). Hormonal effects of several chemicals in recombinant yeast, MCF7 cells and uterotrophic assays in mice. *Microbiological Biotechnology* 10:293-299.

Kang KS, Cho SD, Lee YS (2002). Additive estrogenic activities of the binary mixtures of four estrogenic chemicals in recombinant yeast expressing estrogen receptor. *Journal of Veterinary Science* 3: 1-5.

Lemini C, Silva G, Timossi C, Luque D, Valverde A, Gonzales-Martinez M, Hernandez A, Rubio-Poo C, Chavez Lara B, Valenzuela F (1997). Estrogenic effects of p-hydroxybenzoic acid in CD1 mice. *Environmental Research* 75:130-134.

Nishihara T, Nishikawa JL, Kanayama T, Dakeyama F, Saito K, Imagawa M, Takatori S, Kitagawa Y, Hori S, Utsumi H (2000). Estrogenic activities of 517 chemicals by yeast two-hybrid assay. *Journal of Health Science* 46:282-298.

Okubo T, Yokoyama Y, Kano K, Kano I (2001). ER-dependent estrogenic activity of parabens assessed by proliferation of human breast cancer MCF7 cells and expression of ER alpha and PR. *Food Chemistry and Toxicology* 39:1225-1232.

Pedersen KL, Pedersen SN, Christiansen LB, Korgaard B, Bjerregaard P (2000). The preservatives ethyl-, propyl- and butylparaben are oestrogenic in an in vivo fish assay. *Pharmacology & Toxicology* 86: 110-113.

Routledge EJ, Parker J, Odum J, Ashby J, Sumpter JP (1998). Some alkyl hydroxyl benzoate preservatives (parabens) are estrogenic. *Toxicology and Applied Pharmacology*. 153:12-19.

Soni MG, Gurdock GA, Taylor SL, Greenberg NA (2001). Safety assessment of propyl paraben: A review of the published literature. *Food and Chemical Toxicology* 39:513-532.

Vinagard AM, Kroner W, Lund KH, Bolz U, Peterson JH (2000). Identification and quantification of estrogenic compounds in recycled and virgin paper for household use as determined by an in vitro yeast estrogen screen and chemical analysis. *Chemical Research in Toxicology* 13: 1214-1222.

2. Darbre PD, Aljarrah A, Miller WR, Coldham NG, Sauer MJ, Pope GS (2004). Concentrations of parabens in human breast tumors. *Journal of Applied Toxicology* 24:5-13.

3. Elder RRL (1984). Final report on the safety assessment of methylparaben, ethylparaben, propylparaben and butylparaben. *Journal of the American College of Toxicology* 3:147-209.

4. Darbre PD, Aljarrah A, Miller WR, Coldham NG, Sauer MJ, Pope GS (2004). Concentrations of parabens in human breast tumors. *Journal of Applied Toxicology* 24:5-13.

5. McGrath KG (2003). An earlier age of breast cancer diagnosis related to more frequent use of antiperspirants/deodorants and underarm shaving. *European Journal of Cancer Prevention* 12:479-485.

6. Harvey PW, Darbre P (2004). Endocrine disruptors and human health: Could oestrogenic chemicals in body care cosmetics adversely affect breast cancer incidence in women? A review of evidence and call for further research. *Journal of Applied Toxicology* 24:167-176.

7. Fisher JS, Turner KJ, Brown D, Sharpe RM (1999). Effect of neonatal exposure to estrogenic compounds on development of the ducts of rat testis through puberty to adulthood. *Environmental Health Perspectives* 107:397-405.

Kang KS, Che JH, RYu DY, Kim TW, Li GX, Lee YS (2002). Decreased sperm number and motile activity on the F1 offspring maternally exposed to butyl p-hydroxybenzoic acid (butyl paraben). *Journal of Veterinary Medical Science* 64:227-235.

Oishi S (2002). Effects of propyl paraben on the male reproductive system. Food and Chemical Toxicology 40:1807-1813.

[What We Do](#) | [Events](#) | [Action Center](#) | [Giving](#) | [News](#) | [Resources](#) | [About Us](#) | [Media Center](#) | [Contact Us](#) | [Privacy Policy](#)

© Breast Cancer Fund All Rights Reserved
Breast Cancer Fund
1388 Sutter Street, Suite 400
San Francisco, CA 94109-5400
(415) 346-8223

Copyright 2004